

Bibliografía

- Hirschfield, R. M. A., Keller, M. B., Panico, S., Arons, B. S., Barlow, D., Davidoff, F., et al. (1997). The National Depressive and Manic-Depressive Association consensus statement on the undertreatment of depression. *Journal of the American Medical Association*, 277, 333-340.
- Hollon, S. D., Munoz, R. F., Barlow, D. H., Beardslee, W. R., Bell, C. C., Bernal, G., et al. (2002). Psychosocial intervention development for the prevention and treatment of depression: Promoting innovation and increasing access. *Biological Psychiatry*, 52, 610-630
- Ma, S. H., y Teasdale, J. D. (2004). Mindfulness-based cognitive therapy for depression: Replication and exploration of differential relapse prevention effects. *Journal of Consulting and Clinical Psychology*, 72, 31-40.
- National Institute for Clinical Excellence (2004). *Depression: Management of depression in primary and second care* (Clinical Guideline No. 23). www.nice.org.uk/CG023NICEguideline
- Segal, Z. V., Williams, J. M. G., y Teasdale, J. D. (2002). *Mindfulness-based cognitive therapy for depression: A new approach to preventing relapse*. New York: Guildford Press.
- Teasdale, J. D., Segal, Z. V., Williams, J. M. G., Ridgeway, V. A., Soulsby, J. M., y Lau, M A. (2000). Prevention of relapse/recurrence in major depression by mindfulness-based cognitive therapy. *Journal of Consulting and Clinical Psychology*, 68, 615-623.
- van Schaik, D. J. F., Klein, A. F. J., van Hout, H. P. J., van Marwijk, H. W. J., Beekman, A. T. F., de Haan, M., et al. (2004). Patients' preferences in the treatment of depressive disorder in primary care. *Psychiatry an Primary Care*, 26, 184-189.