

Bibliografía sobre intervención temprana en la psicosis

- Bentall, R.P. y Morrison, A.P. (2002). More harm than good: The case against using antipsychotic drugs to prevent severe mental illness. *Journal of Mental Health, 11*, 351-356.
- Bechdolf, A., Veith,V., Schwarzer, D., Schormann, M., Stamm, E., Janssen, B., Berning, J., Wagner, M. y Klosterkötter, J. (2005). Cognitive-behavioral, a therapy in the pre-psychotic phase: An exploratory study. *Psychiatry Research, 136*, 251-255.
- Birchwood, M., Todd, P. y Jackson, C. (1998). Early intervention in Psychosis: The critical-period hypothesis. *International Clinical Psychopharmacology, 13*, 31-40.
- Cornblatt, B.A. (2002). The New York high risk project to the Hillside recognition and prevention (RAP) program. *American Journal of Medical Genetics, 114*, 956-966.
- Falloon, I.R.H. (1992). Early intervention for first episodes of schizophrenia: A preliminary exploration. *Psychiatry, 55*, 4-14.
- Haddock, G., Lewis, S., Addington, J. y Gleeson, J. (2005). Implementing cognitive-behavioural therapy for first-episode psychosis. *The British Journal of Psychiatry, 187*, 72-76.
- Henry, L. P., McGorry, P.D., Edwards, J., Hulbert, C.A., Henry, L., Harrigan, S., Dudgeon, P., Francey, S., Maude, D., Cocks, J., Killackey, E. y Power, P. (2005). A controlled trial of cognitively oriented psychotherapy for early psychosis (COPE) with four-year follow-up readmission. *Psychological Medicine, 35*, 1295-1306.
- Herrmann-Doig, T., Maude, D. y Edwards, J. (2003). *Systematic Treatment of Persistent Psychosis (STOPP): A Psychological Approach to Facilitating Recovery in Young People with First Episode Psychosis*. Londres: Martin Dunitz.
- McGlashan, T.H. (1999). Duration of untreated psychosis in first episode schizophrenia: marker o determinant of course?. *Biological Psychiatry, 46*, 899-907.
- Morrison, A.P., Bentall, R.P., French, P., Walford, L., Kilcommons, A., Knight, A., Kreutz, M. y Lewis, S.W. (2002). Randomised controlled trial of early detection and cognitive therapy for preventing transition to psychosis in high-risk individuals. *British Journal of Psychiatry, 181*, 78-84.
- Phillips, L (2002). The PACE clinic: identification and management of young people at "Ultra" high risk of psychosis. *Journal of Psychiatric Practice, 8*, 255-269.
- Vallina, O., Alonso, M., Gutiérrez, A., Ortega, J.A., García, A., Fernández, P. y Lemos, S. (2003). Aplicación de un programa de intervención temprana en psicosis. Un nuevo desarrollo para las unidades de salud mental. *Avances en Salud Mental Relacional, 2*.