

Referencias:

Adam, S., Bay, C., Bonsang, E., Germain S. y Perelman, S. (2006). Occupational activities and cognitive reserve: a frontier approach applied to the survey on health, aging and retirement in Europe (SHARE). CREPP Working Paper 2006/05.

Ball, K., Berch, D. B., Helmers, K. F., Jobe, J. B., Leveck, M. D.; Marsiske, M., et al. for the «Advanced Cognitive Training for Independent and Vital Elderly Study Group» (2002). Effects of cognitive training interventions with older adults: a randomized controlled trial». The Journal of the American Medical Association (JAMA) 288, 2271-2281.

Bandura, A. (1997). Self-efficacy: The exercise of control. New York: Freeman.

Barnes, D.E., Alexopoulos, G.S., López, O.L., Williamson, J.D. y Yaffe, K. (2006). Depressive symptoms, vascular disease, and Mild Cognitive Impairment. Findings from the Cardiovascular Health Study. Archives of General Psychiatry 63, 273-279.

Bierman, E.J., Comijs, H.C., Rijmen, F., Jonker, C. y Beekman, A.T. (2008). Anxiety symptoms and cognitive performance in later life: results from the longitudinal aging study Amsterdam. Aging and Mental Health 12, 517-523.

Biessels, G.J., Staekenborg, S., Brunner, E., Brayne, C. y Scheltens, P. (2006) Risk of dementia in diabetes mellitus: a systematic review. Lancet 5: 64-74.

Butler, R.N (1969). Age-ism: another form of bigotry. Gerontologist 9, 243-262.

Craik, F.I.M. (2003). Cambios en la memoria humana relacionados con la edad. En: D. Park y N. Schwarz (Eds.). Envejecimiento cognitivo (pp. 77-93). Madrid: Editorial Médica Panamericana.

Dotson, V.M., Beydoun, M.A. y Zonderman, A.B. (2010). Recurrent depressive symptoms and the incidence of dementia and mild cognitive impairment. Neurology 75, 27-34.

Féart, C., Samieri, C., Rondeau, V., Amieva, H., Portet, F., Dartigues, J.F., et al. (2009). Adherence to a Mediterranean diet, cognitive decline and risk of dementia. The Journal of the American Medical Association (JAMA) 302, 638-648.

Gallacher, J., Bayer, A., Fish, M., Pickering, J., Pedro, S., Dunstan, F., et al. (2009). Does anxiety affect risk of dementia? Findings from the Caerphilly prospective study. Psychosomatic Medicine 71, 659-666.

Geda, Y. E.; Roberts, R.; Knopman, D.; Pankratz, V. S.; Christianson, T.; Boeve, B., et al (2009). Cognitive activities are associated with decreased risk of Mild Cognitive Impairment: The Mayo Clinic Population-Based Study of Aging (estudio presentado en la 61ª Reunión Anual de la Academia Americana de Neurología en Seattle, 25 de abril – 2 de mayo de 2009).

Geda, Y.E., Rosebud, O.R., Knopman, D.S., Christianson, T., Pankratz, V.S., Ivnik, R.J., et al. (2010). Physical exercise, aging, and Mild Cognitive Impairment. A population-based study. Archives of Neurology 67, 80-86.

Giles, H., Mulac, A., Bradac, J.J., y Johnson, P. (1987). Speech accommodation theory: The last decade and beyond. En: M.L. McLaughlin (Ed.). Communication

yearbook 10 (pp. 13-48). Beverly Hills, CA: Sage.

Giles, H., Coupland, N., y Coupland, J. (1991). Accommodation theory: Communication, context and consequence. En: H. Giles, J. Coupland y N. Coupland (Eds.). Contexts of accommodation: Developments in applied linguistics (pp-1-68). Cambridge, UK: Cambridge University Press.

Gramunt (2010). Vive el envejecimiento activo. Barcelona: Obra Social "La Caixa".

Gross, A.L. y Rebok, G.W. (2011). Memory training and strategy use in older adults: Results from the ACTIVE study. *Psychology and Aging*, 28. [Epub ahead of print]

Gross, A.L., Rebok, G.W., Unverzagt, F.W., Willis, S.L. y Brandt, J. (2011). Word list memory predicts everyday function and problema-solving in the elderly: results from the ACTIVE cognitive intervention trial. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn* 18, 129-46.

Gross, A.L., Rebok, G.W., Unverzagt, F.W., Willis, S.L. y Brandt, J. (2011). Cognitive predictors of everyday functioning in older adults: Results from the ACTIVE cognitive intervention trial. *The Journals of Gerontology, Series B: Psychological Sciences and Social Sciences* . May 9. [Epub ahead of print].

Hertzog, C., Kramer, A.F., Wilson, R.S. y Lindenberger, U. (2008). Enrichment effects on adult cognitive development. Can the functional capacity of older adults be preserved and enhanced? *Psychological Science in the Public Interest* 9 (1), 1-65.

Johns Hopkins Medicine. Memory lapses – Normal aging or something more serious? Johns Hopkins Health Alerts, October 6, 2006. www.johnshopkinshealthalerts.com. Con acceso el 15 de mayo de 2011.

Katzman, R., Brown, T., Thal, L.J., Fuld, P.A., Aronson, M., Butters, N., et al. (1988). Clinical, pathological and neurochemical changes in dementia: a subgroup with preserved mental status and numerous neocortical plaques. *Annals of Neurology*, 23, 138-144.

Kivipelto, M., Helkala, E.L., Laakso, M.P., Hänninen, T., Hallikainen, M. y Alhainen, K. (2001) Midlife vascular risk factors and Alzheimer's disease in later life: longitudinal, population based study. *British Medical Journal* 322, b1447-b1451.

Kuypers, J.A., y Bengston, V.L. (1973). Social breakdown and competence: a model of normal aging. *Human Development*, 16, 181-201.

Luchsinger, J.A., Reitz, C., Honig, L.S., Tang, M.X., Shea, S. y Mayeux, R. (2005) Aggregation of vascular risk factors and risk of incident Alzheimer disease. *Neurology* 65: 545-551.

McGuinness, B., Todd, S., Passmore, P. y Bullock, R. (2009). Blood pressure lowering in patients without prior cerebrovascular disease for prevention of cognitive impairment and dementia. *Cochrane Database Syst Rev* 4: CD004034, DOI: 10.1002/14651858.CD004034.pub3.

Middleton, L.E. y Yaffe, K. (2009). Promising strategies for the prevention of dementia. *Archives of Neurology* 66, 1210-1215.

Plassman, B.L., Williams J.W., Burke, J.R., Holsinger, T. y Benjamin, S. (2010).

Systematic review: factors associated with risk for and possible prevention of cognitive decline in later life. *Annals of Internal Medicine*, 153, 182-193.

Qiu, C. y Winblad, B. (2005) The age dependent relation of blood pressure to cognitive function and dementia. *The Lancet Neurology* 4: 487-499.

Reiman, E.M., Chen, K., Langbaum, J.B.S., Lee, W., Reschke, C., Brandy, D., et al. (2010) Higher serum total cholesterol levels in late middle age are associated with glucose hypometabolism in brain regions affected by Alzheimer's disease and normal aging. *Neuroimage* 49: 169-176.

Ritchie, K., Carrière, I., Ritchie, C.W., Berr, C., Artero, S. y Ancelin, M. (2010). Can we design prevention programs to reduce dementia incidence? A prospective study of modifiable risk factors. *British Medical Journal* 5, 341.

Scarmeas, N., Stern, Y., Mayeux, M.D., Manly, J., Schupf, N. et al. (2009). Mediterranean diet and Mild Cognitive Impairment. *Archives of Neurology* 66: 216-225.

Teri, L., Logsdon, R. y McCurry, S. (2008). Exercise interventions for dementia and cognitive impairment: The Seattle protocols. *Journal of Nutrition Health and Aging* 12, 391-394.

Warren, J.D., Schott, J.M., Fox, N.C., Thom, M., Revesz, T., Holton, J.L. et al. (2005). Brain biopsy in dementia. *Brain*, 128, 2016-2025.

Wilson, R.S., Hoganson, G.M., Rajan, K.B., Barnes, L.L., Mendes de Leon, C.F. y Evans, D.A. (2010). Temporal course of depressive symptoms during the development of Alzheimer disease. *Neurology* 75, 21-26.

Wilson, D., Peters, R., Ritchie, K. y Ritchie C.W. (2011). Latest advances on interventions that may prevent, delay or ameliorate dementia. *Therapeutic Advances in Chronic Disease* 2, 161-173.

Whitehouse PJ, George D (2008). The myth of Alzheimer's. What you aren't being told about today's most dreaded diagnosis. New York: St. Martin's Press.

Xu, W., Caracciolo, B., Wang, H.X., Winblad, B., Bäckman, L., Qiu, C. et al. (2010) Accelerated progression from mild cognitive impairment to dementia in people with diabetes. *Diabetes* 59, 2928-2935.

Yaffe, K., Fiocco, A.J., Lindquist, M.S., Vittinghoff, E., Simonsick, E.M., Newman, A.B., et al. (2009). Predictors of maintaining cognitive function in older adults. The Health ABC Study. *Neurology* 72, 2029-2035.